

## SEDATIVE STUDY OF MIDAZOLAM IN GOATS (*CAPRA HIRCUS*)

S. K. JANGRA<sup>1</sup>, S. K. CHAWLA, A. KUMAR, R. TAYAL, JIT SINGH<sup>2</sup>  
and S. M. BEHL

Department of Veterinary Surgery and Radiology  
College of Veterinary Sciences, CCS Haryana Agricultural University, Hisar -125 004

### ABSTRACT

A sedative study was conducted in 5 healthy goats after intravenous administration of midazolam hydrochloride (0.4 mg/kg). The animals went into lateral recumbency by three minutes of midazolam administration. Mild to moderate watery salivation was observed. The palpebral, corneal and swallowing reflexes were diminished after midazolam administration. Relaxation of the jaw, flaccidity of the tongue and relaxation of neck was observed. Limbs, tail and anal sphincters were mildly relaxed. Abdominal muscles relaxation was noticed for about 15 minutes. The head rightening reflex was seen at  $18.2 \pm 3.28$  min after drug administration. The animals stood on their own by  $43.4 \pm 4.47$  min with slight ataxia. Complete recovery occurred after  $69.2 \pm 5.06$  minute.

**Key words:** Midazolam, goat, premedication

Midazolam, an imidazobenzodiazepine derivative, has been used as sedative and induction agent in human beings (Berggren and Eriksson, 1981), dogs (Greene *et al.*, 1993) and pigs (Smith *et al.*, 1991). It has an early induction, short duration of action, rapid elimination and total body clearance time (Court and Greenblatt, 1992). Midazolam is compatible with 5 per cent dextrose in water, 0.9 per cent sodium chloride and lactated Ringer's solution (Brown *et al.*, 1993). Hence, the maintenance of anaesthesia becomes easier when injected along with intravenous fluids. There are sporadic reports of work on midazolam in different species but no systematic work has been conducted in goats. The present study describes the evaluation of sedative effects of midazolam in goats.

### MATERIALS AND METHODS

The present study was conducted on 5 clinically healthy goats (*Capra hircus*) of either sex. All the goats were kept under similar managerial conditions. The goats were controlled in lateral recumbency and midazolam

hydrochloride was administered into the jugular vein at dose rate of 0.4 mg/kg and were immediately let free. Behavioral parameters like spontaneous activity, weak time, down time, relaxation of jaw, abdominal muscles, tongue, neck, tail and limbs, onset of salivation, palpebral/corneal reflex, swallowing reflex, peniculous reflex, sleeping time, time taken to return to sternal recumbency from lateral recumbency and keeping the head upright were recorded. Recovery from the effect of midazolam hydrochloride was judged as time taken by the animal to stand and restoration of co-ordination of limbs.

### RESULTS AND DISCUSSION

The effective intravenous dose of midazolam hydrochloride (0.4 mg/kg.) as a sedative agent was standardized by pilot trials as already found effective for restraining cows (Sanchez *et al.*, 1994). Light sedation was observed with 0.3 mg/kg intravenous administration in human (Kanto *et al.*, 1982). Berggren and Erickson (1981) observed smooth and reliable induction of anaesthesia in human with intravenous dose of  $0.36 \pm 0.01$  mg/kg of midazolam.

Within three minutes of intravenous

<sup>1</sup>Veterinary Surgeon, Department of Animal Husbandry (Haryana Govt.)

<sup>2</sup>Additional Director of Research

