

UTERINE LAVAGE, LEVAMISOLE, CLOPROSTENOL AND ITS COMBINATIONS IMPROVE PLASMA PROTEINS AND FERTILITY OUTCOMES IN ENDOMETRITIC BUFFALOES

NARENDRA SINGH, BHOOPENDRA SINGH¹ and RAJESH KUMAR^{1*}

Department of Veterinary Clinical Complex, R.R. College of Veterinary and Animal Science,
Deoli, Tonk, RAJUVAS, Rajasthan-304 804, India

¹Department of Veterinary Gynaecology & Obstetrics, College of Veterinary Science and AH,
ANDUAT, Ayodhya-224 229 (UP), India

Received: 07.12.2022; Accepted: 28.02.2023

ABSTRACT

The study was aimed to assess comparative efficacy of uterine lavage, levamisole subcutaneously, PGF_{2α} (Cloprostenol) intramuscularly and their combinations on biochemical parameters and treatment outcomes in buffaloes with endometritis. Total thirty-six repeat breeder buffaloes with endometritis were allocated as untreated control (G0, n=6), G1 (Uterine Lavage), G2 (Levamisole, s/c), G3 (Cloprostenol, i/m), G4 (Uterine Lavage+Levamisole, s/c) and G5 (Uterine Lavage + Cloprostenol, i/m), each group comprised of 6 animals. After treatment total protein, albumin and globulin concentration were increased significantly ($p<0.05$) in treated buffaloes. The highest clinical cure was recorded with uterine lavage plus levamisole (83.33%) and uterine lavage plus Cloprostenol (83.33%) treated buffaloes followed by G2 (66.67%) and G3 (66.67%), G1 (50.00%) and G0 (33.33%) group. The highest conception rates was in group G4 and G5 (66.67% each) followed by G2 and G3 (50% each), G1 (33.00%) and G0 (16.67%). In conclusion, endometritic buffaloes can be treated using lavage of uterus with physiological saline in combination with parenteral Levamisole or Cloprostenol.

Keywords: Buffalo, Cloprostenol, Endometritis, Levamisole, Uterine lavage

How to cite: Singh, N., Singh, B. and Kumar, R. (2023). Uterine lavage, levamisole, cloprostenol and its combinations improve plasma proteins and fertility outcomes in endometritic buffaloes. *The Haryana Veterinarian* 62(SI-2): 117-119.

Subclinical endometritic and non-specific infections are most common cause of repeat breeding syndrome in buffaloes. Clinical endometritis is defined as purulent vaginal discharge (21 days or more postpartum) or a muco-purulent vaginal discharge (after 26 days postpartum). while subclinical endometritis is characterized by absence of muco-purulent or purulent vaginal discharge (Sheldon *et al.*, 2006) of uterine origin. Subclinical endometritis is often undiagnosed (Dutt *et al.*, 2017), hence remain untreated and results in subfertility. Transrectal and ultrasonographic examination of uterus are usually performed to diagnose the disease under field conditions (Barlund *et al.*, 2008).

The therapeutic regimen used to treat infected bovine uterus should effectively clear uterine infection without compromising its local immunity and adulterating the milk or meat. Many therapeutic agents have been tried and tested in treatment of infected uterus (Singh *et al.*, 2018; Ahmadi *et al.*, 2019). In view of development of antimicrobial resistance and various side effects with use of antibiotics; the levamisole may be used to substitute or supplement antibiotic therapeutics (Singh *et al.*, 2017). In equines, the infection of uterus can be treated with physiological saline solution as uterine lavage. The uterine lavage results in removal of bacteria, non-functional neutrophils, hasten myometrial contractility which in turns

causes evacuation of uterine content and influx of fresh neutrophils in uterine lumen (Brinsko *et al.*, 2011). Thus, the present work was planned with the hypothesis that lavage of uterus along with administration of Levamisole or Cloprostenol parenterally, would cure subclinical endometritis and improve reproductive outcomes of such buffaloes. Thus, this study was carried out to assess aforesaid approaches on plasma proteins and treatment outcomes in repeat breeder buffaloes with endometritis.

MATERIALS AND METHODS

Referral cases presented at Veterinary Clinical Complex, C.V.Sc. & A.H, ANDUAT, Ayodhya, were screened for study following approval from ethical committee of university (No. IAEC/CVSc/2019/P-02). Based on animal breeding history and records, trans-rectal palpation of genitalia and White side test, total 36 repeat breeder buffaloes were assigned in six groups, each comprised of 6 buffaloes viz. G0 (untreated control), G1: Uterine lavage (UL) with 300 ml NSS in divided doses, 50 mL each time, as slow IU infusion), G2: Levamisole (LEV) @ 2.5 mg/kg body weight subcutaneously, three occasions i.e. 0, 7th and 14th day of the cycle; G3: Cloprostenol or CP @ 250 µg intramuscularly, on day-10 of the cycle; G4: (G1+G2 combined i.e. UL+LEV) and G5 (G1+G3 combined i.e. UL+CP). Six repeat breeder buffaloes were served as untreated control. For biochemical analysis, 6 ml of blood was collected via jugular vein puncture in

*Corresponding author: drrajesh25@gmail.com

clean sterile vials and centrifuged at 1200xg for 15 min at 4 °C. Thereafter, the plasma was separated and preserved at -20 °C until analysis. The plasma was analyzed for estimation of total protein, albumin, globulin, using diagnostic kits as per manufacturer instructions. Blood sample was collected twice, first at commencement of experiment (BT, before therapy) and second at consequent estrus following treatment (AT, after therapy). The double insemination was done at 12 h interval using frozen thawed semen in estrus buffaloes. The animals were inseminated again at second and third subsequent estrus, if returned to estrus. In order to diagnose pregnancy, trans-rectal palpation was done at 45-60 days after AI. The data were analyzed using Duncan's MRT and complete randomized design at significance of $p < 0.05$ level (Snedecor and Cochran, 1994).

RESULTS AND DISCUSSION

TOTAL PROTEIN: The total protein (gm/dl) (mean $\pm$ SE) elevated significantly in treated buffaloes (G1 to G5) than their corresponding values before treatment, however in untreated animals, the values were did not differ significantly (Table 1). Similar to present observations, lower serum protein levels were recorded in endometritic buffaloes (Patil, 2010) and cows (Sivaraman *et al.*, 2003) but there is also evidence to contrary findings (Ahmad *et al.*, 2004; Sharma, 2018). The significant increase in the serum protein level of endometritic buffaloes following therapy with UL might be due to dilution of bacteria and elimination of micro organisms, debris and products of uterine inflammation that improved general body condition of the animal and thereby increased serum protein level in the animal after treatment. Our findings are corroborated well with earlier studies (Biswal *et al.*, 2014; Singh *et al.*, 2017). To date, no literature cited regarding effect of CP on serum protein level in endometritic buffaloes. So, our results could not be compared. The significant increase in the serum protein level of endometritic buffaloes after treatment with UL+LEV and UL+CP, might be due to combined effect of uterine lavage with LEV/CP. Scanning of literature revealed that no citation available concerning effect of uterine UL+LEV and uterine UL+CP on serum protein level of endometritic buffaloes. So, our results could not be compared.

ALBUMIN: The Albumin (gm/dl)(mean $\pm$ SE) levels were elevated significantly in treated buffaloes (G1 to G5) than their corresponding pretreatment values, whereas the values did not differ significantly among untreated control buffaloes (Table 1). Like present findings, earlier workers also recorded lower level of serum albumin in endometritic cattle (Sivaraman *et al.*, 2003; Amle *et al.*, 2014) and buffaloes (Patil, 2010; Sharma, 2018). In present study, intrauterine infusion of saline may cause dilution of infection and elimination of microorganisms,

debris and inflammatory exudates from the uterus that improved general body condition of the animal and thereby increased serum albumin level in the animal after treatment. Our findings are consistent with earlier reports on UL and LEV subjected endometritic animals (Patil, 2010; Biswal *et al.*, 2014; Singh *et al.*, 2017). The significant increase in the serum albumin level of endometritic buffaloes after treatment with uterine UL+LEV and UL+CP might be due to combined effect of uterine lavage and levamisole. Scanning of literature revealed that no citation available concerning effect of Cloprostenol, uterine UL+CP and UL+LEV on serum albumin level in endometritic buffaloes. So, our results could not be compared.

The significant increase in the serum albumin level of endometritic buffaloes after treatment with levamisole might be due to its immunomodulatory effect. The concentration of albumin is negatively correlated with acute phase proteins i.e. inflammation and infection results in declined albumin level (Eckersall and Bell, 2010) and elevation of albumin concentration following levamisole therapy was might be due to its immune-potentiating properties. The significant increase in the serum albumin level of endometritic buffaloes after treatment with CP might be due to its luteolytic effect (which in turns results in declined progesterone concentration and hence improved uterine immunity), stimulation of myometrial contractility, phagocytic activity of uterine polymorphonuclear, hence improved general body condition of the buffaloes thereby increased plasma albumin level of animals.

GLOBULIN: The Globulin level (gm/dl) (mean $\pm$ SE) elevated significantly in treated buffaloes (G1 to G5) than their corresponding values before treatment, whereas the values did not differ significantly among control buffaloes (Table 1). Similar to our observations, lower serum globulin levels were reported in endometritic cattle (Sivaraman *et al.*, 2003; Magnus and Lali (2009) and buffaloes (Patil, 2010). Our findings are consistent with earlier reports on UL and LEV treated endometritic animals (Patil, 2010; Singh *et al.*, 2017). The significant increase in the serum globulin level of endometritic buffaloes following therapy with lavage of uterus with normal saline might be due to elimination of uterine debris, microorganisms, dilution of inflammatory contents as well as infection, which in turns results in improved body condition of the animal and thereby increased serum globulin level in the animal after treatment. Furthermore, the significant increase in the serum globulin level of endometritic buffaloes after treatment with levamisole might be due to immunomodulatory property of levamisole. The significant increase in the serum globulin level of endometritic buffaloes after treatment with Cloprostenol CP might be due to its luteolytic effect (which

Table 1. Effect of different treatment on plasma proteins and fertility outcomes in endometritic buffaloes

Groups (n=6)	Total Protein (g/dL)		Albumin (g/dL)		Globulin (g/dL)		Fertility outcomes	
	BT	AT	BT	AT	BT	AT	RR	CR
G0 (Untreated)	6.07±0.05	6.16±0.05 ^D	2.74±0.02	2.81±0.03 ^C	3.33±0.06	3.35±0.05 ^C	33.33 ^D (2/6)	16.67 ^D (1/6)
G1 (UL)	6.19±0.08 ^a	7.39±0.08 ^{Cb}	2.68±0.04 ^a	2.91±0.07 ^{Bb}	3.51±0.09 ^a	4.48±0.11 ^{Bb}	50.00 ^C (3/6)	33.33 ^C (2/6)
G2 (LEV)	6.18±0.09 ^a	8.10±0.08 ^{Ab}	2.66±0.04 ^a	3.02±0.03 ^{Ab}	3.52±0.06 ^a	5.07±0.07 ^{Ab}	66.67 ^B (4/6)	50.00 ^B (3/6)
G3 (CP)	6.17±0.07 ^a	7.90±0.05 ^{Bb}	2.62±0.04 ^a	2.95±0.06 ^{Bb}	3.53±0.05 ^a	4.95±0.06 ^{Ab}	66.67 ^B (4/6)	50.00 ^B (3/6)
G4 (UL+LEV)	6.13±0.07 ^a	8.20±0.05 ^{Ab}	2.64±0.03 ^a	3.12±0.03 ^{Ab}	3.48±0.09 ^a	5.08±0.10 ^{Ab}	83.33 ^A (5/6)	66.67 ^A (4/6)
G5 (UL+CP)	6.20±0.06 ^a	8.16±0.06 ^{Ab}	2.69±0.04 ^a	3.09±0.05 ^{Ab}	3.50±0.09 ^a	5.07±0.09 ^{Ab}	83.33 ^A (5/6)	66.67 ^A (4/6)

Means with different superscripts within group (a, b) and between groups (A, B, C) differ significantly ($p < 0.05$); BT (Before therapy); AT (After therapy); RR (Response rate); CR (Conception rate).

in turns results in declined progesterone concentration and hence improved uterine immunity), stimulation of myometrial contractility, phagocytic activity of uterine polymorphonuclear, hence improved general body condition of the buffaloes thereby increased serum globulin level of animals. The significant increase in the serum globulin level of endometritic buffaloes after treatment with uterine lavage plus Levamisole and uterine lavage plus Cloprostenol might be due to combined effect of uterine lavage and levamisole. To date, no literature is cited regarding effect of Cloprostenol, Lavage of uterus with saline plus Levamisole and Lavage of uterus with saline plus Cloprostenol on serum globulin level in endometritic buffaloes. So, our results could not be compared.

CONCLUSION

In conclusion, Levamisole and Cloprostenol alone were found to be more effective than the uterine lavage therapy. The higher cure rate was recorded in uterine lavage along with parenteral Levamisole or Cloprostenol treated buffaloes as compared to any regimen alone; as evidenced by maximum improvement in plasma protein level, recovery rate and conception rate was observed in levamisole plus uterine lavage and Cloprostenol plus uterine lavage group. The sub-clinical endometritis in buffaloes can be treated using Levamisole (@ 2.5 mg/kg body weight via s/c route on three occasions at 7 days interval), furthermore, it can be used as a substitute or supplement to antibiotic therapy.

REFERENCES

Ahmad, I., Lodhi, L.A., Qureshi, Z.I. and Younis, M. (2004). Studies on blood glucose, total proteins, urea and cholesterol levels in cyclic, non-cyclic and endometritic crossbred cows. *Pak. Vet. J.* **24**(2): 92-94.

Ahmadi, M.R., Makki, M., Mirzaei, A. and Gheisari, H.R. (2019). Effects of hypertonic dextrose and paraffin solution as non-antibiotic treatments of clinical endometritis on reproductive performance of high producing dairy cows. *Reprod. Domest. Anim.* **54**(5): 762-771.

Amle, M., Padodkar, V.P., Shehlar, R. and Birade, H. (2014). Serum biochemical levels of repeat breeder cross bred cows under rural condition of Satara district Maharashtra. *Int. J. Adv. Vet. Sci. Tech.*

3(1): 109-113.

Barlund, C.S., Carruthers, T.D., Waldner, C.L. and Palmer, C.W. (2008). A comparison of diagnostic techniques for postpartum endometritis in dairy cattle. *Theriogenol.* **69**: 714-723.

Biswal, S., Das, S., Mohanty, D.N. and Jena, D. (2014). Effect of systemic and local immunomodulation therapies on conception rate in endometritic cows. *Indian J. Field Vet.* **10**(2): 1-4.

Brinsko, S.P., Blanchard, T.L., Varner, D.D., Schumacher, J., Love, C.C., Hinrichs, K. and Hartman, D.L. (2011). Endometritis. *Manual Equine Reprod.* pp. 73-84.

Dutt, R., Singh, G., Singh, M., Sharma, M., Dalal, J. and Chandolia, R.K. (2017). Diagnosis of subclinical endometritis in Murrah buffaloes through Cytobrush technique. *Int. J. Curr. Microbiol. Appl. Sci.* **6**(11): 494-499.

Eckersall, P.D. and Bell, R. (2010). Acute phase proteins: Biomarkers of infection and inflammation in veterinary medicine. *Vet. J.* **185**: 23-27.

Magnus, P. and Lali, K.F. (2009) Serum Biochemical Profile of Post-Partum Metritic Cow. *Vet. World* **2**(1): 27-28.

Patil, R.D. (2010). Studies on haemato-biochemical, hormonal and therapeutic aspects of postpartum metritis in Surti buffaloes. M.V.Sc thesis submitted to Anand Agricultural University, Gujarat (India).

Sharma (2018). Therapeutic efficacy of ceftiofur sodium, gentamicin sulphate and zingibar officinale extract on recovery of endometritic buffaloes. M.V.Sc thesis submitted to Narendra Deva University of Agriculture and Technology, Ayodhya, Uttar Pradesh.

Sheldon, I.M., Lewis, G.S., LeBlanc, S. and Gilbert, R.O. (2006). Defining post-partum uterine disease in cattle. *Theriogenol.* **65**: 1516-1530.

Singh, B., Gupta, H.P., Prasad, S. and Rajora, V.S. (2018). Effect of Immunomodulators on serum nitric oxide concentration of endometritic repeat breeding crossbred cows. *Indian J. Vet. Res.* **27**(1): 32-35.

Singh, P.P., Pande, N., Bhavna and Agrawal, R. (2017). Clinical and biochemical studies on endometritic repeat breeding cows following treatment with levamisole. *The Haryana Veterinarian* **56**(1): 55-57.

Sivaraman, T., Shanmugasundaram, S., Arunachalam, S. and Sivakumar, T. (2003). Blood profile constituents associated with production diseases in Jersey crossbred cows. *Indian J. Anim. Sci.* **73**(1): 44-47.

Snedecor, G.W. and Cochran, W.G. (1994). Statistical methods. (8th Edn.), Iowa State University Press, Ames, Iowa, USA.